

**GCE**

**Physics A**

**H556/01: Modelling physics**

Advanced GCE

**Mark Scheme for Autumn 2021**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.



This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2021

Annotations available in RM Assessor

	Annotation	Meaning
	Correct response	Used to indicate the point at which a mark has been awarded ( <b>one tick per mark awarded</b> ).
	Incorrect response	Used to indicate an incorrect answer or a point where a mark is lost.
<b>AE</b>	Arithmetic error	Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
<b>BOD</b>	Benefit of doubt given	Used to indicate a mark awarded where the candidate provides an answer that is not totally satisfactory, but the examiner feels that sufficient work has been done.
<b>BP</b>	Blank page	Use BP on additional page(s) to show that there is no additional work provided by the candidates.
<b>CON</b>	Contradiction	No mark can be awarded if the candidate contradicts himself or herself in the same response.
<b>ECF</b>	Error carried forward	Used in <u>numerical answers only</u> , unless specified otherwise in the mark scheme. Answers to later sections of numerical questions may be awarded up to full credit provided they are consistent with earlier incorrect answers. Within a question, ECF can be given for AE, TE and POT errors but not for XP.
<b>L1</b>	Level 1	L1 is used to show 2 marks awarded and L1 <sup>^</sup> is used to show 1 mark awarded.
<b>L2</b>	Level 2	L2 is used to show 4 marks awarded and L2 <sup>^</sup> is used to show 3 marks awarded.
<b>L3</b>	Level 3	L3 is used to show 6 marks awarded and L3 <sup>^</sup> is used to show 5 marks awarded.
<b>POT</b>	Power of 10 error	This is usually linked to conversion of SI prefixes. Do not allow the mark where the error occurs. Then follow through the working/calculation giving ECF for subsequent marks if there are no further errors.
<b>SEEN</b>	Seen	To indicate working/text has been seen by the examiner.
<b>SF</b>	Error in number of significant figures	Where more SFs are given than is justified by the question, do not penalise. Fewer significant figures than necessary will be considered within the mark scheme. <b>Penalised only once in the paper.</b>
<b>TE</b>	Transcription error	This error is when there is incorrect transcription of the correct data from the question, graphical read-off, formulae booklet or a previous answer. Do not allow the relevant mark and then follow through the working giving ECF for subsequent marks.
<b>XP</b>	Wrong physics or equation	Used in <u>numerical answers only</u> , unless otherwise specified in the mark scheme. Use of an incorrect equation is wrong physics even if it happens to lead to the correct answer.
<b>^</b>	Omission	Used to indicate where more is needed for a mark to be awarded (what is written is not wrong but not enough).

Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

<b>Annotation</b>	<b>Meaning</b>
/	alternative and acceptable answers for the same marking point
<b>Reject</b>	Answers which are not worthy of credit
<b>Not</b>	Answers which are not worthy of credit
<b>Ignore</b>	Statements which are irrelevant
<b>Allow</b>	Answers that can be accepted
( )	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
<b>ECF</b>	Error carried forward
<b>AW</b>	Alternative wording
<b>ORA</b>	Or reverse argument

## SECTION A

Question	Answer	Marks	Guidance
1	C	1	
2	A	1	
3	C	1	
4	B	1	
5	A	1	
6	B	1	
7	B	1	
8	D	1	
9	D	1	
10	C	1	
11	C	1	
12	B	1	
13	C	1	
14	B	1	
15	A	1	
	<b>Total</b>	<b>15</b>	

## SECTION B

**General rule:** For substitution into an equation, allow any subject – unless stated otherwise in the guidance

Question			Answer	Marks	Guidance
16	(a)	(i)	$(g \rightarrow) [\text{m s}^{-2}]$ and $(t \rightarrow) [\text{s}]$ or $(gt^2 \rightarrow) [\text{m s}^{-2} \times \text{s}^2]$ Clear evidence of working leading to m on both sides	<b>M1</b> <b>A1</b>	
		(ii)	s / distance measured with a ruler / tape measure Timer mentioned for measuring t / time Measure distance from bottom of ball to (top of) trapdoor Any <u>one</u> from: <ul style="list-style-type: none"> <li>Take repeated readings (for t for same s) to determine average t</li> <li>Avoid parallax error when using the ruler</li> </ul>	<b>B1</b> <b>B1</b> <b>B1</b> <b>B1</b>	
	(b)	(i)	$(p_1 = 4.4 \times 0.050) = 0.22 \text{ (kg m s}^{-1}\text{)}$	<b>B1</b>	
		(ii)	(impulse =) $\frac{1}{2} \times 30 \times 0.02$ or $0.30 \text{ (kg m s}^{-1}\text{)}$  $-0.30 = p_2 - 0.22$  $p_2 = (-) 0.08 \text{ (kg m s}^{-1}\text{)}$	<b>C1</b> <b>C1</b> <b>A1</b>	Allow any correct re-arrangement Possible ECF from <b>(b)(i)</b>  <b>Ignore sign</b> <b>Allow 0.52 for 2 marks</b>
		(iii)	(momentum of trapdoor =) $0.30 \text{ (kg m s}^{-1}\text{)}$  $v = 3.0 \text{ (m s}^{-1}\text{)}$	<b>C1</b> <b>A1</b>	Allow $(KE_{\text{trapdoor}} =) \frac{1}{2} \times 0.05 \times (4.4^2 - 1.6^2)$ or $0.42 \text{ (J)}$ Possible ECF from <b>(b)(ii)</b>  <b>Allow 1 SF answer here</b> <b>Allow alternate methods involving CoE (giving 2.9) and e(giving 2.8)</b>
<b>Total</b>				<b>12</b>	

Question		Answer	Marks	Guidance
17	(a)	(mean) = 1.87(2) (mm) (range) = 0.04 mm (percentage uncertainty =) $\frac{0.02}{1.872} \times 100$ percentage uncertainty = 1 (%)	<b>C1</b>  <b>C1</b>  <b>A1</b>	<b>Allow</b> use of resolution of micrometer (gives percentage uncertainty of 0.5%) <b>Allow</b> use of maximum or minimum deviation from the mean  <b>Allow</b> 2 or 3 SF answer

	(b)*	<p><b>Level 3 (5–6 marks)</b> Clear description <b>and</b> clear analysis <i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p><b>Level 2 (3–4 marks)</b> Some description <b>and</b> some analysis <b>or</b> Clear description <b>or</b> Clear analysis <i>There is a line of reasoning presented with some structure. The information presented is in the most-part relevant and supported by some evidence.</i></p> <p><b>Level 1 (1–2 marks)</b> Limited description <b>or</b> Limited analysis <i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b> No response or no response worthy of credit.</p>	<b>B1× 6</b>	<p>Use level of response annotation in RM Assessor.</p> <p><b>Indicative scientific points may include:</b></p> <p><b>Description</b></p> <ul style="list-style-type: none"> <li>• Determine <math>T</math> by measuring several oscillations</li> <li>• Independent and dependent variables identified (e.g. <math>L</math> and <math>T</math>)</li> <li>• Variables kept constant (e.g. for <math>L</math> and <math>T</math> experiment, <math>m</math> is kept constant)</li> <li>• Repeating to determine average <math>T</math></li> <li>• Measure length <math>L</math> and width <math>w</math> with ruler</li> <li>• Measure thickness <math>t</math> with a vernier (calliper) / micrometer</li> <li>• Use video/phone camera / stopwatch / data-logger and motion sensor / light gates and timer</li> <li>• Use top-pan balance / scales to measure <math>m</math></li> </ul> <p><b>Analysis</b></p> <ul style="list-style-type: none"> <li>• Plot an <u>appropriate</u> graph, e.g. <math>T^2</math> against <math>L^3</math> <b>or</b> tabulate <math>T^2 \div L^3</math></li> <li>• Gradient of <b>best line</b> determined <b>or</b> average of <math>T^2 \div L^3</math></li> <li>• Use a large triangle to determine gradient</li> <li>• Gradient (or equivalent) related to <math>E</math>, e.g. gradient = <math>16\pi^2 m / wEt^3</math> for <math>T^2</math> against <math>L^3</math> graph</li> </ul>
		<b>Total</b>	<b>9</b>	



Question			Answer	Marks	Guidance
18	(a)	(i)	(energy =) $150 \times 7$ <b>or</b> 1050 (J)  $1050 = 0.025 \times c \times 20$  (c =) $2100 \text{ (J kg}^{-1} \text{ K}^{-1}\text{)}$	<b>C1</b>  <b>C1</b>  <b>A1</b>	<b>Allow</b> any correct re-arrangement
		(ii)	(energy=) $150 \times (63 - 7)$ <b>or</b> 8400 (J)  $8400 = L_{(f)} \times 0.025$  ( $L_f =$ ) $3.4 \times 10^5 \text{ (J kg}^{-1}\text{)}$	<b>C1</b>  <b>A1</b>	
		(iii)	Longer time to heat water (through the same temperature) / shorter time to heat (ice) through same temperature / gradient of graph is greater for ice / gradient of graph is smaller for water/AW  Water has greater specific heat capacity	<b>M1</b>  <b>A1</b>	
(b)	(i)	Molecules in <b>X</b> vibrate about fixed positions /AW  Molecules in <b>Z</b> are free to move/random/AW	<b>B1</b>  <b>B1</b>	<b>Allow</b> references to ice for <b>X</b> and water / liquid for <b>Z</b>  <b>Allow</b> <u>one</u> correct for <b>B1</b> from: <ul style="list-style-type: none"> <li>• Molecules in <b>X</b> have lower <u>KE</u>/speed/velocity</li> <li>• Speed/velocity of molecules increases with temp/time</li> <li>• Amplitude or frequency increases with temp/time in <b>X</b></li> </ul>	

		(ii)	<table border="1"> <thead> <tr> <th>Region</th> <th>Physical quantity, or quantities, that increase as time increases</th> <th>Physical quantity, or quantities, that remain constant as time increases</th> </tr> </thead> <tbody> <tr> <td>X</td> <td>K</td> <td>P</td> </tr> <tr> <td>Y</td> <td>P</td> <td>K</td> </tr> <tr> <td>Z</td> <td>K</td> <td>P</td> </tr> </tbody> </table>	Region	Physical quantity, or quantities, that increase as time increases	Physical quantity, or quantities, that remain constant as time increases	X	K	P	Y	P	K	Z	K	P	B1×3	Note that each B1 mark is for a correct row Allow KP/- for both X and Z
Region	Physical quantity, or quantities, that increase as time increases	Physical quantity, or quantities, that remain constant as time increases															
X	K	P															
Y	P	K															
Z	K	P															
		(iii)	Absolute zero / 0 <u>K</u> / - 273 <u>°C</u>	B1													
			<b>Total</b>	<b>13</b>													

Question		Answer	Marks	Guidance
19	(a)	For a body in (rotational) equilibrium the sum of the clockwise moments (about any point) is equal to the sum of the anticlockwise moments (about the same point)	<b>B1</b>	<b>Note</b> Accept $\Sigma$ / total (AW) for sum
	(b)	(i)		
		(i)	(horizontal component of $F$ ) $F \times \cos 20^\circ$  $F \cos 20^\circ \times 1.30 = 0.30 \times 40 \times 9.81$  $F = 96.4$ (N)	<b>M1</b>  <b>M1</b>  <b>A1</b> <b>Allow</b> ECF for incorrect trig i.e. use of sine (gives $F = 265$ ) or $\cos(20 \text{ radians})$ which gives $F = 222$ for 2 marks. <b>Allow ECF</b> for incorrect units for angle and incorrect trig $\sin(20 \text{ radians})$ which gives $F = 99(.2)$ for 1 mark
		(ii)		
		(ii)	$R = F \cos 20^\circ$ or $96(.4) \times \cos 20^\circ$  ( $R =$ ) 91 (N)	<b>C1</b> <b>Allow</b> ECF from (b)(i)  <b>A1</b> <b>Answer is 90.6 (N) to 3sf if 96.4 used.</b> <b>Answer is 90(.2) (N) to 3sf if 96 used</b>
	(c)	When (line of force of the) weight falls to the right of the (bottom of the) wheel/AW	<b>B1</b>	
<b>Total</b>			<b>7</b>	

Question			Answer	Marks	Guidance
20	(a)	(i)	Straight symmetrical radial field lines and correct direction of field	B1	Ignore field lines inside the Earth
		(ii)	X and Y labelled which should be an equal distance away from the centre of the Earth	B1	Note Judge by eye Allow X and Y both on the surface of the Earth

	(b)*	<p><b>Level 3 (5–6 marks)</b> Clear description <b>and</b> correct calculations leading to value of total energy (must include the negative sign) <i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p><b>Level 2 (3–4 marks)</b> Some description and some correct calculations <b>or</b> Correct calculations (including the negative sign) <i>There is a line of reasoning presented with some structure. The information presented is in the most-part relevant and supported by some evidence.</i></p> <p><b>Level 1 (1–2 marks)</b> Limited description <b>or</b> Limited calculations <i>The information is basic and communicated in an unstructured way. The information is supported by limited evidence and the relationship to the evidence may not be clear.</i></p> <p><b>0 marks</b> No response or no response worthy of credit.</p>	B1×6	<p><b>Indicative scientific points may include:</b></p> <p><b>Description</b></p> <ul style="list-style-type: none"> <li>• Orbit above the equator / equatorial orbit</li> <li>• Orbit from west to east/same direction of orbit as Earth's rotation</li> <li>• Orbital period is 24 hours / 1 (sidereal) day /23hrs 56mins (4 s)</li> <li>• Orbit is circular / above the same point on the Earth</li> </ul> <p><b>Calculation</b></p> <ul style="list-style-type: none"> <li>• <math>E = (-)\frac{GMm}{r}</math></li> <li>• <math>E = \frac{6.67 \times 10^{-11} \times 5.97 \times 10^{24} \times 2500}{4.22 \times 10^7} = (-) 2.4 \times 10^{10} \text{ J}</math></li> <li>• <math>v = \frac{2\pi r}{T} = \omega r</math></li> <li>• <math>v = \frac{2\pi \times 4.22 \times 10^7}{24 \times 3600} = 3.07 \times 10^3 \text{ m s}^{-1}</math></li> <li>• <math>E = \frac{1}{2}mv^2</math></li> <li>• <math>E = \frac{1}{2} \times 2500 \times [3.07 \times 10^3]^2 = 1.2 \times 10^{10} \text{ J}</math></li> <li>• Total energy = <math>- 2.4 \times 10^{10} + 1.2 \times 10^{10} = - 1.2 \times 10^{10} \text{ J}</math></li> <li>• Allow full credit for algebraic proof using <math>\frac{GMm}{r^2} = \frac{mv^2}{r}</math>, <math>E = (-)\frac{GMm}{r}</math>, <math>E = \frac{1}{2}mv^2</math> and total energy = KE + PE</li> </ul> <p>Allow higher order answers in terms of Lagrange's Identity</p>
		<b>Total</b>	<b>8</b>	

Question		Answer	Marks	Guidance
21	(a)	$\omega^2 = k/m$ or $60/0.080$ or $\omega^2 = 750$ $T = 2\pi/27.39$ or $T = 0.2295$ (s) $t = \frac{1}{4} \times 0.2295$ $t = 0.057$ (s)	<b>C1</b> <b>C1</b> <b>C1</b> <b>A1</b>	<b>Allow</b> correct algebraic expression for T <b>Allow</b> incorrect value for omega <b>Allow</b> incorrect value of T
	(b) (i)	$E = \frac{1}{2}kx^2$ or $E = mgh$ or $0.080 \times 9.81 \times 0.20$ or $\frac{1}{2} \times 60 \times x^2$ $0.080 \times 9.81 \times 0.20 = \frac{1}{2} \times 60 \times x^2$ $x = 0.072$ (m)	<b>C1</b>  <b>C1</b> <b>A1</b>	
	(ii)1	Time of flight is independent of speed/AW Because distance of fall is the same <b>and</b> initial velocity vertically is zero / velocity is horizontal at <b>X</b>	<b>B1</b>  <b>B1</b>	<b>Allow</b> algebraic answers that assume initial vertical velocity is zero/velocity is horizontal at X.
	(ii)2	$D$ increases as speed at <b>X</b> increases because the time of flight is constant/AW $D$ is directly proportional to speed at <b>X</b>	<b>M1</b>  <b>A1</b>	<b>Allow</b> $d = vt$ idea “ $D$ is directly proportional to speed at <b>X</b> because the time of flight is constant” scores 2.
<b>Total</b>			<b>11</b>	

Question			Answer	Marks	Guidance
22	(a)	(i)	$\{v = \omega r \text{ and } \omega = 2\pi f\}$ or $v = 2\pi fr$  Comparison with $y = mx$ leading to gradient = $2\pi r$ or $\Delta v / \Delta f = 2\pi r$	B1  B1	Allow $v/f = 2\pi r$
		(ii)	Line of best fit drawn  Gradient = 62.5 (m)  $2\pi r = 62.5$  $r = 9.9$ (m)	B1  M1  M1  A0	Allow $\pm 3$  Allow ECF on gradient
		(iii)	$F = \frac{mv^2}{r}$ or $F = ma$ and $a = \frac{v^2}{r}$  $F = \frac{1.7 \times 10^{-27} \times [2.0 \times 10^7]^2}{9.9}$  $F = 6.8 \times 10^{-14}$ (N)	C1  C1  A1	Allow use of candidate's answer for (ii) or use of '10'  Expect answers of 6.8 or $6.9 \times 10^{-14}$ (N)
	(b)		$r \propto v^2$ / speed increases by a factor of $\sqrt{2}$  maximum speed = $2.8 \times 10^7$ (m s <sup>-1</sup> )	C1  A1	Allow substitution into correct equation with r doubled  Allow recalculation from previous value of force in (a)(iii)
			<b>Total</b>	<b>10</b>	

Question		Answer	Marks	Guidance	
23	(a)	$g = \frac{G \times 6.31 \times 10^{30}}{(1.90 \times 10^9)^2}$ $g = 117 \text{ (N kg}^{-1}\text{)}$ $2 \times \frac{0.14}{1.90} + \frac{0.42}{6.31} \quad \text{or} \quad 0.21 \quad \text{or} \quad 21\%$ <p>(absolute uncertainty =) 25 (N kg<sup>-1</sup>)</p>	<p><b>C1</b></p> <p><b>C1</b></p> <p><b>C1</b></p> <p><b>A1</b></p>	<p><b>Reject</b> 0.42/(0.14<sup>2</sup>) = 21.4</p> <p><b>Note:</b> final answer of <math>g = 117 \pm 25 \text{ (N kg}^{-1}\text{)}</math> with no working scores all 4 marks.</p> <p><b>Allow</b> correct identification of %(r) and %(m) for 1 mark max if no other marks scored.</p> <p><b>Allow</b> alternate method using max/min values of m and r that give the correct absolute uncertainty (28)</p>	
	(b)	(i)	The (total radiant) power (of a star) /AW	<b>B1</b>	
		(ii)	$(L = 4\pi r^2 \sigma T^4)$ $\text{ratio} = \sqrt{\left(\frac{6.92 \times 7500^4}{10.0 \times 4500^4}\right)}$ $\text{ratio} = 2.3(1)$	<p><b>C1</b></p> <p><b>A1</b></p>	<p><b>Allow</b> 1 mark for 5.3; square root omitted</p> <p><b>Allow</b> 1 mark for 1:2.3 or 0.43</p>
		(iii)	$\lambda_{(\text{max})} \propto \frac{1}{T}$ <p>Lower temperature star will have the longest wavelength, so it is Aa2</p>	<p><b>B1</b></p> <p><b>B1</b></p>	<p><b>Allow</b> word equation</p> <p><b>Note</b> Must mention wavelength <u>and</u> temperature</p>
		(iv)	From their (different) colours	<b>B1</b>	



	(v)	<p>Any <u>three</u> from:</p> <ul style="list-style-type: none"> <li>• Continuous spectrum</li> <li>• Light / radiation / photons passes through cooler gas/star's atmosphere</li> <li>• Photon(s) absorbed by electron(s)</li> <li>• Electron(s) excited / jump / make transition to higher energy level(s)</li> <li>• Electron only promoted if energy of photon matches energy gap between two given levels</li> <li>• Photons remitted in different directions</li> <li>• (so) idea of contrast with non-absorbed wavelengths</li> </ul>	B1 × 3	
	(c)	<p>Any <u>two</u> from:</p> <ul style="list-style-type: none"> <li>• Black hole has smaller mass / radius / size</li> <li>• Black hole has higher density/gravitational field strength/stronger gravitational field</li> <li>• black hole absorbs light / does not emit visible light</li> <li>• Has an escape velocity =&gt; c</li> <li>• No fusion in a black hole (ORA)</li> </ul>	<p>B1</p> <p>B1</p>	<p><b>Allow</b> black hole emits Hawking radiation</p>
		<b>Total</b>	<b>15</b>	

**OCR (Oxford Cambridge and RSA Examinations)**  
**The Triangle Building**  
**Shaftesbury Road**  
**Cambridge**  
**CB2 8EA**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

[www.ocr.org.uk](http://www.ocr.org.uk)

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored