



A Level Physics Online

AQA Physics - 7407/7408

Module 9: Astrophysics

You should be able to demonstrate and show your understanding of:	Progress and understanding:			
	1	2	3	4
9.1 Telescopes				
9.1.1 Astronomical telescope consisting of two converging lenses				
Ray diagram to show the image formation in normal adjustment.				
Angular magnification in normal adjustment.				
$M = \text{angle subtended by image at eye} / \text{angle subtended by object at unaided eye}$				
Focal lengths of the lenses.				
$M = f_o / f_e$				
9.1.2 Reflecting telescopes				
Cassegrain arrangement using a parabolic concave primary mirror and convex secondary mirror.				
Ray diagram to show path of rays through the telescope up to the eyepiece.				
Relative merits of reflectors and refractors including a qualitative treatment of spherical and				
9.1.3 Single dish radio telescopes, I-R, U-V and X-ray telescopes				
Similarities and differences of radio telescopes compared to optical telescopes. Discussion should include structure, positioning and use, together with comparisons of resolving and collecting powers.				
9.1.4 Advantages of large diameter telescopes				
Minimum angular resolution of telescope.				
Rayleigh criterion:				
$\theta \approx \lambda / D$				
Comparison of the eye and CCD as detectors in terms of quantum efficiency, resolution and convenience of use.				



You should be able to demonstrate and show your understanding of:	Progress and understanding:			
	1	2	3	4
No knowledge of the structure of the CCD is required.				
9.2 Classification of stars				
9.2.1 Classification by luminosity				
Apparent magnitude, m .				
The Hipparcos scale.				
Dimmest visible stars have a magnitude of 6.				
Relation between brightness and apparent magnitude. Difference of 1 on magnitude scale is equal to an intensity ratio of 2.51.				
Brightness is a subjective scale of measurement.				
9.2.2 Classification by luminosity				
Parsec and light year.				
Definition of M , relation to m : $m - M = 5 \log (d / 10)$				
9.2.3 Classification by temperature, black-body radiation				
Stefan's law and Wien's displacement law.				
General shape of black-body curves, use of Wien's displacement law to estimate black-body temperature of sources.				
Experimental verification is not required.				
$\lambda_{max}T = constant = 2.9 \times 10^{-3} m K$				
Assumption that a star is a black body.				
Inverse square law, assumptions in its application.				
Use of Stefan's law to compare the power output, temperature and size of stars $P = \sigma AT^4$				



You should be able to demonstrate and show your understanding of:				Progress and understanding:																																				
				1	2	3	4																																	
9.2.4 Principles of the use of stellar spectral classes																																								
<table border="1"> <thead> <tr> <th>Spectral class</th> <th>Intrinsic colour</th> <th>Temperature / K</th> <th>Prominent absorption lines</th> </tr> </thead> <tbody> <tr> <td>O</td> <td>blue</td> <td>25 000 – 50 000</td> <td>He⁺, He, H</td> </tr> <tr> <td>B</td> <td>blue</td> <td>11 000 – 25 000</td> <td>He, H</td> </tr> <tr> <td>A</td> <td>blue-white</td> <td>7 500 – 11 000</td> <td>H (strongest) ionized metals</td> </tr> <tr> <td>F</td> <td>white</td> <td>6 000 – 7 500</td> <td>ionized metals</td> </tr> <tr> <td>G</td> <td>yellow-white</td> <td>5 000 – 6 000</td> <td>ionized & neutral metals</td> </tr> <tr> <td>K</td> <td>orange</td> <td>3 500 – 5 000</td> <td>neutral metals</td> </tr> <tr> <td>M</td> <td>red</td> <td>< 3 500</td> <td>neutral atoms, TiO</td> </tr> </tbody> </table>				Spectral class	Intrinsic colour	Temperature / K	Prominent absorption lines	O	blue	25 000 – 50 000	He ⁺ , He, H	B	blue	11 000 – 25 000	He, H	A	blue-white	7 500 – 11 000	H (strongest) ionized metals	F	white	6 000 – 7 500	ionized metals	G	yellow-white	5 000 – 6 000	ionized & neutral metals	K	orange	3 500 – 5 000	neutral metals	M	red	< 3 500	neutral atoms, TiO					
Spectral class	Intrinsic colour	Temperature / K	Prominent absorption lines																																					
O	blue	25 000 – 50 000	He ⁺ , He, H																																					
B	blue	11 000 – 25 000	He, H																																					
A	blue-white	7 500 – 11 000	H (strongest) ionized metals																																					
F	white	6 000 – 7 500	ionized metals																																					
G	yellow-white	5 000 – 6 000	ionized & neutral metals																																					
K	orange	3 500 – 5 000	neutral metals																																					
M	red	< 3 500	neutral atoms, TiO																																					
Description of the main classes.																																								
Temperature related to absorption spectra limited to Hydrogen Balmer absorption lines: requirement for atoms in an $n = 2$ state.																																								
9.2.5 The Hertzsprung-Russell (HR) diagram																																								
General shape: main sequence, dwarfs and giants.																																								
Axis scales range from -10 to $+15$ (absolute magnitude) and 50 000 K to 2500 K (temperature) or OBAFGKM (spectral class).																																								
Students should be familiar with the position of the Sun on the HR diagram.																																								
Stellar evolution: path of a star similar to our Sun on the HR diagram from formation to white dwarf.																																								
9.2.6 Supernovae, neutron stars and black holes																																								
Defining properties: rapid increase in absolute magnitude of supernovae; composition and density of neutron stars; escape velocity $> c$ for black holes.																																								
Gamma ray bursts due to the collapse of supergiant stars to form neutron stars or black holes.																																								
Comparison of energy output with total energy output of the Sun.																																								
Use of type 1a supernovae as standard candles to determine distances. Controversy concerning accelerating Universe and dark energy.																																								
Students should be familiar with the light curve of typical type 1a supernovae.																																								



You should be able to demonstrate and show your understanding of:	Progress and understanding:			
	1	2	3	4
Supermassive black holes at the centre of galaxies.				
Calculation of the radius of the event horizon for a black hole, Schwarzschild radius (R_S) $R_S \approx 2GM / c^2$				
9.3 Cosmology				
9.3.1 Doppler effect				
$\Delta f / f = v / c$ and $z = \Delta \lambda / \lambda = -v / c$ for $v \ll c$ applied to optical and radio frequencies.				
Calculations on binary stars viewed in the plane of orbit.				
Galaxies and quasars.				
9.3.2 Hubble's law				
Red shift $v = Hd$				
Simple interpretation as expansion of universe; estimation of age of universe, assuming H is constant.				
Qualitative treatment of Big Bang theory including evidence from cosmological microwave background radiation, and relative abundance of hydrogen and helium.				
9.3.3 Quasars				
Quasars as the most distant measurable objects.				
Discovery of quasars as bright radio sources.				
Quasars show large optical red shifts; estimation involving distance and power output.				
Formation of quasars from active supermassive black holes.				
9.3.4 Detection of exoplanets				
Difficulties in the direct detection of exoplanets.				
Detection techniques will be limited to variation in Doppler shift (radial velocity method) and the transit method.				
Typical light curve.				

